



## Effect of the Natural Extract of *Juglans Mandshurica* Epicarp on Broiler Coccidiosis Disease

Kyongho Choe<sup>1</sup> , Myongdok Im<sup>1</sup> , Jeman Chae<sup>1</sup> , Cholmin Kim<sup>2</sup> , Sunil Choe<sup>3</sup> , Myongil Jin<sup>3</sup> ,  
Chunsik Ri<sup>1</sup>  & Unju Son<sup>1</sup> 

<sup>1</sup> Pyongyang University of Agriculture, Department of Veterinary Medicine and Animal Husbandry, Pyongyang 999093, DPR of Korea

<sup>2</sup> Bioengineering Branch, State Academy of Sciences, Pyongyang 999093, DPR of Korea

<sup>3</sup> Kye Ung Sang Sariwon University of Agriculture, Department of Veterinary Medicine and Animal Husbandry, Sariwon, 950003, DPR of Korea

Poultry Science Journal 2023, 11(2): 257-268

### Keywords

ACI  
Extract  
Broiler  
Coccidiosis  
Plant medicine

### Corresponding author

Kyongho Choe  
cuijinghao1981@163.com

### Article history

Received: March 25, 2023  
Revised: June 06, 2023  
Accepted: August 21, 2023

### Abstract

Coccidiosis is a protozoal disease caused by *Eimeria* that significantly impacts the global poultry industry. In this study, *in vitro* and *in vivo* experiments were conducted to evaluate the anticoccidial effect of *Juglans mandshurica* epicarp aqueous extract as an alternative treatment against coccidiosis in broiler chickens. In the *in vitro* experiment, unsporulated *Eimeria tenella* oocysts were exposed to various concentrations of the aqueous extract. After 48 hours of incubation, the degree of inhibition of sporulation and morphed oocysts were examined. For the *in vivo* experiment, seven groups of two-week-old broiler chickens were divided into A1-A5, positive control (PC), and negative control (NC) groups. The A1-A4 groups received different doses of the aqueous extract calculated based on toxicity tests, while group A5 was fed diets mixed with salinomycin. Groups A1-A5 were infected with *E. tenella*, while PC was infected but unmedicated, and NC was uninfected and unmedicated. Results showed that the aqueous extract inhibited the sporulation of *E. tenella* oocysts *in vitro* in a concentration-dependent manner. All experimental groups fed with the extract exhibited significantly higher weight gain, particularly those receiving 5 mL/kg BW. Moreover, the groups receiving doses of 5 and 7 mL/kg BW showed significant differences in bloody Diarrhea and cecal lesion scores compared to the PC group. The anticoccidial index (ACI) value for these groups was above 160, indicating high efficacy similar to the group fed with mixed salinomycin. In conclusion, the study suggests that *J. mandshurica* epicarp aqueous extract could be a safe and effective alternative treatment for coccidiosis in broilers. This could provide a new approach to control coccidiosis in the global poultry industry.

### Introduction

Coccidiosis is one of the main parasitic diseases caused by protozoan *Eimeria* in poultry and is recognized as a serious parasitic disease worldwide (Kostadinović *et al.*, 2015). Seven coccidiosis species have been identified in chicken, among which *Eimeria tenella*, *Eimeria brunetti*, and *Eimeria necatrix* are known to be highly pathogenic (Hafez, 2008). Coccidiosis is a major parasitic disease affecting poultry and results in severe economic loss due to severe reductions in feed utilization and body weight gain (Kostadinović *et al.*, 2015).

Coccidiosis control relies mainly on chemicals, which costs much money. The continuous emergence of antibiotic-resistant *Eimeria* has raised concerns about the safety of anticoccidial medicines and their potential negative impact on human, animal and environmental health (Primož and Suzana, 2012; Nilsson *et al.*, 2012). In this context, the use of anticoccidial chemicals for commercial poultry production is prohibited in many countries and new countermeasures are urgently required along with public distrust of meat using chemical medicines (Giannenas *et al.*, 2003; Applegate *et al.*, 2010; Peek

and Landman, 2011; Bozkurt *et al.*, 2012). It has been strongly advocated that synthetic medicines should be opposed to natural additives (Alaeldein *et al.*, 2018; Rashed and Alaeldein, 2019; Saleh *et al.*, 2020). The use of plant medicament as one of the natural additives is a new idea since it has several advantages including low cost, safety, low residue or no residue in broilers and many other benefits (Kostadinović *et al.*, 2015; Mahmood *et al.*, 2020).

Juglans mandshurica Maxim, also called Manchurian walnut or Onigurumi, is a perennial deciduous broad-leaf tree that is fast-growing in northeastern China and Korean Peninsula (Wei-Ning *et al.*, 2010; Wen-Ting *et al.*, 2016). The bark, branches, epicarp of immature fruits, leaves, roots, and stems have high therapeutic effects and have thus been extensively used to cure a variety of diseases in the past (Yan *et al.*, 2019). For instance, the epicarp of immature fruits has been demonstrated to be against cancer, dermatosis, diarrhea, dysentery, gastric ulcer, leukopenia, and uterine prolapse (Qing *et al.*, 2010; Arvind *et al.*, 2011; Yuanyuan *et al.*, 2015; Yuwei *et al.*, 2016; Jin-Hai *et al.*, 2018; Yuan-Yuan *et al.*, 2019). The significant anti-inflammatory activity has been reported for extracts of *J. mandshurica* in several inflammatory models (Xiaohui *et al.*, 2015; Fei *et al.*, 2021). In mice, the treatment with the juglone in the extracts of the immature exocarps of *J. mandshurica* significantly decreased the levels of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6, proving that it suppressed the inflammatory response through inhibition of the TLR4/NF- $\kappa$ B signaling pathway (Xiaohui *et al.*, 2015).

Juglone (5-hydroxy-1, 4-naphthoquinone), a type of naphthoquinone, is a phenolic compound that exhibits allelopathic properties. It has remarkable antibacterial activity against Gram-positive, Gram-negative bacteria, yeast and fungi (Saling *et al.*, 2011; Qin *et al.*, 2019; Fei *et al.*, 2021).

However, to the best of our knowledge, no studies have been published on the use of *J. mandshurica* epicarp as an anticoccidial agent. In addition, since *J. mandshurica* epicarp has been regarded as a discarded part not used for food, its beneficial use in the poultry industry has not yet been explored.

In this study, we have assumed that natural plant extracts can be used as alternatives to synthetic coccidiosis drugs. Therefore, we tested the anticoccidial effect of the aqueous extract of *J. mandshurica* epicarp.

## Materials and Methods

All the animal experiments in our paper were performed by the guidelines of the Institutional Animal Care and Use Committee of DPR of Korea and approved by the Ethics Committee of the Institute of veterinary medicine of the Academy of Agricultural Researchers. All efforts

were made to minimize suffering.

### Preparation of Eimeria oocyst samples

A single species of *E. tenella* oocyst was isolated from infected chicken caecum by saltwater float selection and subjected to specific modifications. Unsporulated oocysts were obtained from the caecal contents of chickens 7 days after infection. The oocysts in saturated saline solution were centrifuged for 10 min at 1000g at 4°C, and the supernatant containing the oocysts was transferred to a 1L measuring cylinder filled with distilled water. The cylinder was left overnight at 4 °C and then the upper solution was removed not to disturb the pellets, leaving the final volume of 100 mL. Under the abovementioned conditions, the remaining solution (100mL) was further centrifuged to remove the supernatant, and 10 mL of the solution containing oocysts was obtained. Oocysts size was calculated from samples of eighty oocysts isolated from the chicken caeca. The morphology and quantity of oocysts were analyzed under the microscope and finally confirmed by PCR (Fernandez *et al.*, 2003). The oocysts were extracted with a DNA extraction kit (Takara, Japan), and PCR was performed with a Taq DNA polymerases kit (Takara, Japan). The PCR conditions are as follows; an initial denaturation at 94°C for 3 min, 30 cycles of 45 sec at 94 °C, 30 sec at 60°C, and 1 min at 72 °C for annealing, with a final extension step at 72 °C for 5 min. The amplification products were separated on 1.5% agarose gel followed by ethidium bromide staining and visualized under UV light.

To enhance sporulation, the oocysts were put in incubator glass dishes containing 2.5% potassium dichromate solution, which was then aspirated and stirred on the occasion. Daily examination of the dishes proceeded until 90–95% sporulation. The dishes were washed using phosphate buffer solution (PBS, pH=7.2) 3 times before use and diluted to the desired concentration of 1 mL solution.

### Preparation of plant extracts and medicine feeding

*J. mandshurica* epicarp was collected in the mountains around Pyongyang city, DPR of Korea, between August and September. The epicarp of *J. mandshurica* was dried at 37 °C for 48 hours and extracted with water (volume ratio of 1:10) at 37 °C for 2 days. Table 1 shows representative active ingredients of *J. mandshurica* epicarp extract.

The chickens were fed a medicated mixture of feed and medication, where the quantity of medication was calculated based on their body weight. Following the medicated feed, they were given their regular feed and water. The medication was administered during the first feeding of the day for seven consecutive days (Giannenas *et al.*, 2014).

Table 1. The composition of the active ingredients of the extract

No	Chemical constituents	Extracts	Parts
1	Juglone	Water	Green walnut husks
2	2-Hydroxy-1,4-naphthoquinone	Water	Green walnut husks
3	Phenolics	Water	Green walnut husks
4	Tetralones	Water	Green walnut husks
5	Triterpenoids	Water	Green walnut husks
6	Flavonoids	Water	Green walnut husks
7	Phenylpropanoids	Water	Green walnut husks
8	Coumarins	Water	Green walnut husks
9	Steroids	Water	Green walnut husks

**In vitro examination procedure**

Non-sporulated *E. tenella* oocysts were exposed to the aqueous extract of different concentrations (0.01, 0.1, 1, 10, 30, 60, 100%) of *J. mandshurica* epicarp. After incubation at room temperature (26-29°C) for 48 hours, the number of sporulated, non-sporulated, and deformed oocysts was recorded as a percentage. The NC group was exposed to distilled water, and all the experiments were repeated three times.

**In vivo examination procedure**

A total of two-week-old Cobb chicks (n = 350) (Institute of veterinary medicine of the Academy of Agriculture, Pyongyang, DPR of Korea) were acclimated for one week before the experiment. The test groups consisted of 5 groups (A1-A5, n=10 per group) and the chickens were orally administered with  $4 \times 10^4$  sporulated oocysts (Table 2).

Table 2. In vivo experimental design<sup>1</sup>

Groups	Chickens	Applied medicines	Dosage mL/kg BW	Days of medicine application
NC	10			
PC	10			
A1	10	The aqueousextract of <i>J. mandshurica</i> epicarp	1	7
A2	10	The aqueousextract of <i>J. mandshurica</i> epicarp	3	7
A3	10	The aqueousextract of <i>J. mandshurica</i> epicarp	5	7
A4	10	The aqueousextract of <i>J. mandshurica</i> epicarp	7	7
A5	10	Salinomycin	66 mg/kg	7

<sup>1</sup>Five replicates were performed for each group of the experiment.

PC: positive control (infected without medication); NC: negative control (uninfected without medication).

Table 3. Composition and nutritional composition of standard diet<sup>1</sup>

Ingredient	Amount, %
Corn	58.2
Soybean meal	28.4
Corn gluten meal	8.4
Corn oil	2.0
Dicalcium phosphate	1.4
Salt	0.3
DL-Methionine	0.2
Lysine	0.1
Vitamin and trace element premix <sup>2</sup>	1.0
Calculated nutrient composition	Amount
ME, kcal/kg	2,830
Crude Protein, %	19.03
Methionine+Cysteine, %	0.9
Methionine, %	0.47
Lysine, %	1.05
Calcium, %	0.95
Available phosphorus, %	0.43
Total phosphorus, %	0.60

<sup>1</sup>Experimental chicks were fed a diet based on the breeder's nutrient recommendations for the Cobb 700 breeder (Cobb-Vantress, 2020)

<sup>2</sup>Minimum guarantee analysis of vitamin and trace element premix: Vit A 10000 IU/kg; Vit D3 1500 IU/kg; Vit E 5 mg/kg; Vit K3 1 mg/kg; Vit B1 0.5 mg/kg; Vit B2 2 mg/kg; Vit B5 (pantothenic acid) 5 mg/kg; Vit B6 5 mg/kg; Vit B12 0.2 mg/kg; Vit PP 10 mg/kg; B9 (folic acid) 0.5 mg/kg; biotin 1 mg/kg; choline chloride 500mg; Fe 60 mg; Zn 60 mg; Mn 60 mg; Cu 5.6 mg; I 0.5 mg; Co 0.3 mg.

The medicine doses of the A1-A4 groups were calculated at 1, 3, 5, and 7 mL/kg BW respectively. The medicine was applied daily from the beginning of oral oocysts administration. The A5 group was fed diets containing salinomycin (Institute of veterinary medicine of the Academy of Agriculture, Pyongyang,

### Medicine toxicity test

The toxicity test of *J. mandshurica* epicarp can be divided into acute toxicity test and chronic toxicity test. For the acute toxicity test, mice were divided into five groups (at doses of 4, 5, 6, 7, and 8 g Kg<sup>-1</sup>, respectively) with eight mice in each group. The concentrate of epicarp aqueous extraction was dissolved in a reasonable amount of saline and then administered by intraperitoneal injection. After injection, the number of deaths was recorded for 14 days and LD50 was calculated using the Bliss method. For the chronic toxicity test, two-week-old Cobb chicks were divided into the control and test groups (30 chicks in each group), and the number of deaths, appetite loss, depression index, and body weight change were investigated while feeding the medicine at a therapeutic dose (aqueous extraction of 6ml/kg body weight) for one month.

DPR of Korea) 66mg/kg, while the PC group was infected and without medication, and the NC group was uninfected and without medication. Chickens were reared in wire-floor cages. Feed and water are free to eat and the basic rations are guaranteed by the instructions (Table 3).

### Assessment of anticoccidial activity

The anticoccidial activity of the aqueous extract of *J. mandshurica* epicarp was assessed by bloody diarrhea, relative body weight gain (BWG), survival rate, oocyst output/g excreta, and lesion scores. The anticoccidial index was calculated using Dexing *et al.* (2011) method with the following formula:

ACI (anticoccidial index) = (relative ratio of BWG + survival rate) - (lesion scores + oocyst value)

According to ACI values, the anticoccidial effects were determined as follows; ACI >180 was considered excellent, 180-160 was considered marked, 160-140 was considered moderate, 140-120 was considered slight, and ACI <120 was considered inactive (Yasuhiro *et al.*, 1977).

On the 4th–7th days after infection, the chickens were examined for bloody diarrhea by enumerating bloody excreta twice daily. The degree of bloody diarrhea is the average value of blood excreta described by Abbas *et al.* (2010), which can be divided into 0-3 categories according to the blood content of 0, 33, 33-66, and 66-99% in the fecal. Feces from each group were collected between the 4th and 7th days after infection and the oocysts were counted by a hemocytometer according to Holdsworth *et al.* (2004). One gram of each well-mixed feces sample was

soaked in 10 mL of water, and 10µL of suspension was taken to calculate the oocysts under the optical microscope. The results have been expressed as oocysts/g output. The oocyst value and oocyst reduction rate were calculated using a method described by Lan *et al.* (2016), Li-yun *et al.* (2021), and Min *et al.* (2019).

$$\text{ROP (relative oocyst production)\%} = \frac{\text{Oocysts per g output of every group}}{\text{Oocysts per g output of the PC group}} \times 100$$

When the ROP is 0-1%, the oocyst value is 0; when the ROP is 1-25%, the oocyst value is 5; when the ROP is 26-50%, the oocyst value is 10; when the ROP is 51-75%, the oocyst value was 20; when the ROP was 76-100%, the oocyst value was 40.

$$\text{Oocysts reduction rate(\%)} = \frac{\text{OPG of PC group} - \text{OPG of experimental group}}{\text{OPG of PC group}} \times 100$$

$$\text{Survival rate(\%)} = \frac{\text{Number of surviving chickens}}{\text{initial number of chickens}} \times 100$$

The relative weight gain rate was calculated by measuring the bodyweight of each group from the beginning date of infection to the final date of infection and subtracting the initial BW from the final BW.

$$\text{Relative ratio BWG} = \frac{\text{Average BWG in each group}}{\text{Average BWG of NC group}} \times 100$$

On the seventh day of infection, 5 chickens of each group were slaughtered, and the cecum of each bird was removed and observed. Lesion scores were marked from 0 to 4 depending on the severity of the cecum.

Lesion value = mean lesion score × 10.

The degree of the lesion was determined using the method described by Johnson and Reid (1970). After the initial examination of the appearance of the lesion, the chicken cecum was further cut open to observe the internal lesions of the intestine. According to the severity of chicken cecal lesions, the lesion score was divided into five grades.

### Statistical Analysis

All the data were analyzed by SPSS 20.0 software. Significance of the differences in BWG and oocyst discharge of the NC, PC, A1-5 groups were statistically analyzed by one-way ANOVA and recorded as Mean and SEM. In addition, the nonparametric Kruskal-Wallis H test was used to compare the bloody diarrhea score and lesion value among the NC, PC, A1-5 groups. And Dunn's multiple comparison tests were used to analyze statistical

differences between the NC, PC, A1-5 groups. The significance level was set at  $P < 0.05$ .


## Results

### Morphometric analysis of Eimeria species

Based on morphology, the length and the width of oocysts were determined as  $21.3 \pm 1.44 \mu\text{M}$  and  $18.2 \pm 1.68 \mu\text{M}$  respectively, and the shape index was  $1.17 \pm 1.08$ . The results showed that the isolated oocysts were *E. tenella*. Finally, the oocysts were identified by PCR and the result showed a definite band at 539bp (Figure 1).

### Acute and chronic toxicity tests

In the acute toxicity test, LD50 of the aqueous extract of *J. mandshurica* epicarp was 5.76 g/kg, and its confidence limit was 5.39-6.11g/Kg. The results of chronic toxicity tests demonstrated that there was no difference in the number of deaths, loss of appetite,




**Figure 1.** Agarose gel electrophoresis of PCR products using DNA samples of *E. tenella* (lane 2) and a control with no starting DNA (lane 1). Molecular size markers (2000DNA Marker) in base pairs are indicated on the left.

As shown in Figure 2, at the concentration of 100, 60, 30, 10, and 1% of aqueous extract, the sporulation rate was 16.3, 23.0, 31.7, 46.0, 61.7%, respectively, compared to the water control group (98%), displaying significant sporulation inhibition effect ( $P < 0.05$ ).

and depression when 6mL of the aqueous extract of *J. mandshurica* epicarp (/kg body weight) was given to healthy 2-week-old Cobb chicks every day for one month. However, in terms of weight change, the test groups were slightly higher than the control groups, but there was no significant difference. The results revealed that the extract from the epicarp of *J. mandshurica* had no negative effect on the growth of chicks.


### Sporulation inhibition test

The sporulation inhibition effect of the aqueous extract of *J. mandshurica* epicarp was evaluated by treating isolated *E. tenella* oocyst with various concentrations. In this experiment, the sporulation rate of oocysts was calculated by recording the number of sporulated, non-sporulated, and deformed oocysts at different concentrations of the aqueous extracts (Figure 2).



**Figure 2.** Effect of *J. mandshurica* extract on sporulation of *E. tenella*. *E. tenella* showed sporulation differences in water and different concentration of the aqueous extract of *J. mandshurica* epicarp. Water was used as a negative control (NC). The mean for three replicates is shown, a–d Different letters indicate statistically significant differences ( $P < 0.05$ ).

Very low concentrations of aqueous extract (0.01% and 0.1%) had no significant sporulation inhibition activity. After being incubated with different concentrations of the aqueous extract for 48 hours, more deformed oocysts were detected (Figure 3).



**Figure 3.** Effect of *J. mandshurica* extract on deformation of *E. tenella* oocysts. Water was used as a negative control (NC). The mean for three replicates are shown, a–d Different letters indicate statistically significant differences ( $P < 0.05$ ).

As shown in Figure 3, the aqueous extract of *J. mandshurica* epicarp had a great effect on oocyst deformation. The inhibition effect was especially significant when the concentration of the aqueous extract was 10, 30, 60, and 100% ( $P < 0.05$ ).

#### Effect on relative body weight gain rate

The relative body weight gain rates of chicks infected with *E. tenella* were measured for different doses of aqueous extract (Table 4 and Figure 4).

**Table 4.** Comparison of relative body weight gain, bloody diarrhea, lesion value, oocyst value, and ACI in the different chicken groups

Group	Survival rate (%)	Relative weight gain rate (%)	Bloody diarrhea (Mean)	Lesion value (Mean)	Oocyst value	ACI (Mean)
NC	100.00	100.00 <sup>a</sup>	0.00	0.00	0.00	200.00
PC	80.00	67.40 <sup>d</sup>	3.0 <sup>a</sup>	35 <sup>a</sup>	40.00	72.40
A1	100.00	74.34 <sup>cd</sup>	2.3 <sup>b</sup>	28 <sup>b</sup>	20.00	126.34
A2	100.00	83.01 <sup>c</sup>	1.6 <sup>c</sup>	19 <sup>c</sup>	10.00	154.01
A3	100.00	87.99 <sup>bc</sup>	0.7 <sup>d</sup>	17 <sup>d</sup>	10.00	160.99
A4	100.00	87.40 <sup>bc</sup>	0.6 <sup>d</sup>	17 <sup>d</sup>	10.00	160.40
A5	100.00	90.78 <sup>b</sup>	0.4 <sup>d</sup>	16 <sup>d</sup>	10.00	164.78
SEM	0.84	1.28	0.12	1.22	1.42	2.73
P-Value	<0.0522	<0.038	<0.0269	<0.0285	<0.0313	<0.0116


A1–A4: groups applied with aqueous extract of *J. mandshurica* epicarp; A1: 1 mL aqueous extract/kg bodyweight; A2: 3 mL aqueous extract/kg bodyweight; A3: 5 mL aqueous extract/kg bodyweight; A4: 7 mL aqueous extract/kg bodyweight; A5: 66 mg salinomycin/kg in feed; PC: positive control (infected without medication); NC: negative control (uninfected without medication); ACI: anticoccidial index; a–d Different letters indicate statistically significant differences ( $P < 0.05$ ).

As shown in Figure 4, the relative body weight gain in the experimental groups was significantly different from that of the control group infected with oocysts but did not receive the medicines ( $P < 0.05$ ). Also, among the experimental groups fed with aqueous extract, the relative body weight gain of group A3 fed with 5

mL/kg BW was relatively high.

#### Oocysts per gram of feces (OPG)

No oocysts appeared in the feces within 3 days after oocysts inoculation, and there was a difference in the average excretion of oocysts from 4th to 7th days (Table 5).




**Figure 4.** Relative weight gain of chicks infected with *E. tenella*. A1-A4: groups applied with aqueous extract of *J. mandshurica* epicarp; A1: 1mL aqueous extract/kg bodyweight; A2: 3mL aqueous extract/kg bodyweight; A3: 5mL aqueous extract/kg bodyweight; A4: 7mL aqueous extract/kg bodyweight; A5: 66mg salinomycin/kg in feed; PC: positive control (infected without medication); NC: negative control (uninfected without medication). a–d Different letters indicate statistically significant differences ( $P < 0.05$ ).

**Table 5.** Comparison of mean OPG output calculated in the feces of different groups and inhibition rate

Group	OPG (Mean $\times$ 105)	Inhibition rate (%)
NC	0.00	100.00 <sup>a</sup>
PC	31.62 <sup>a</sup>	0.00 <sup>e</sup>
A1	19.80 <sup>b</sup>	37.38 <sup>d</sup>
A2	12.98 <sup>c</sup>	58.95 <sup>c</sup>
A3	10.74 <sup>d</sup>	66.03 <sup>b</sup>
A4	9.72 <sup>d</sup>	69.26 <sup>b</sup>
A5	8.29 <sup>d</sup>	73.78 <sup>b</sup>
SEM	1.14	3.61
<i>P-Value</i>	<0.034	<0.012

A1-A4: groups applied with aqueous extract of *J. mandshurica* epicarp; A1: 1mL aqueous extract/kg bodyweight; A2: 3mL aqueous extract/kg bodyweight; A3: 5 mL aqueous extract/kg bodyweight; A4: 7 mL aqueous extract/kg bodyweight; A5: 66mg salinomycin/kg in feed; PC: positive control (infected without medication); NC: negative control (uninfected without medication); OPG: oocysts in per gram feces; a–e Different letters indicate statistically significant differences ( $P < 0.05$ ).




**Figure 5.** Effect of *J. mandshurica* extract on oocysts production and reduction rate of *E. tenella* oocysts after oral inoculation. A1-A4: groups applied with aqueous extract of *J. mandshurica* epicarp; A1: 1mL aqueous extract/kg bodyweight; A2: 3mL aqueous extract/kg bodyweight; A3: 5mL aqueous extract/kg bodyweight; A4: 7mL aqueous extract/kg bodyweight; A5: 66mg salinomycin/kg in feed; PC: positive control (infected without medication); NC: negative control (uninfected without medication). a–e Different letters indicate statistically significant differences ( $P < 0.05$ ).

A1-A4 test groups (administered with different doses of the aqueous extract) and the A5 test group (fed with salinomycin mixed food) had significant differences with PC group in oocysts excretion and oocyst reduction rate ( $P < 0.05$ ) (Figure 5).

### Bloody diarrhea and cecal lesion value

All chicks infected with *E. tenella* oocysts developed

bloody diarrhea, but the symptoms in the groups treated with medicine were lighter than those in the PC group. Especially, A5 group and A2, A3, and A4 groups were higher than that of the PC group ( $P < 0.05$ ) (Table. 4 and Figure 6A). In addition, the cecal lesions status of the medicine feeding test group was significantly different from that of the PC group (Table 4 and Figure 6B).




**Figure 6.** Effect of *J. mandshurica* extract on bloody diarrhea and cecal lesion value in chicks after oral infection with *E. tenella* oocysts. A1-A4: groups applied with aqueous extract of *J. mandshurica* epicarp; A1: 1 mL aqueous extract/kg bodyweight; A2: 3 mL aqueous extract/kg bodyweight; A3: 5 mL aqueous extract/kg bodyweight; A4: 7 mL aqueous extract/kg bodyweight; A5: 66 mg salinomycin/kg in feed; PC: positive control (infected without medication); NC: negative control (uninfected without medication). a–d Different letters indicate statistically significant differences ( $P < 0.05$ ).

These results demonstrated that the aqueous extract of *J. mandshurica* epicarp has a positive effect on alleviating bloody Diarrhea and the pathological symptoms caused by coccidiosis.

### Anticoccidial index

In comparison with the PC group, the test groups fed

medicine showed a significant anticoccidial effect, especially the anticoccidial index of groups A3, A4, and A5 was 160 or higher, indicating that the anticoccidial effects were marked according to Yasuhiro *et al.* (1977), (Table. 4 and Figure 7).



**Figure 7.** Effect of *J. mandshurica* extract on the anticoccidial index in chicks after oral infection with *E. tenella* oocysts. A1-A4: groups applied with aqueous extract of *J. mandshurica* epicarp; A1: 1 mL aqueous extract/kg bodyweight; A2: 3 mL aqueous extract/kg bodyweight; A3: 5 mL aqueous extract/kg bodyweight; A4: 7 mL aqueous extract/kg bodyweight; A5: 66 mg salinomycin/kg in feed; PC: positive control (infected without medication); NC: negative control (uninfected without medication). These results illustrated that the aqueous extract of *J. mandshurica* epicarp has a good anticoccidial activity similar to salinomycin.



## Discussion

With the increase in the use of veterinary drugs, the safety issue of livestock products has become a major concern. In particular, human health issues owing to the chemical drug residues in livestock products and reduced therapeutic effects due to drug resistance (Takele, 2016) are recognized as major challenges for the veterinary sector. As one of the ways to solve this problem, plant medicine development is very important in meeting people's demand for natural food and reducing the cost of medicine use (Nahed *et al.*, 2022; Tuhinadri and Samir, 2015). Many plant-derived drugs are currently being used in veterinary medicine, particularly as anticoccidial agents. Several plant-based anticoccidial agents have been introduced, such as *Curcuma longa* L (Abbas *et al.*, 2010), *Rumex nervosus* (Saleh *et al.*, 2020), *Pinus radiata* (Abdul *et al.*, 2009), and *Brucea javanica* (Lan *et al.*, 2016). These plant-based veterinary drugs are not only effective as anticoccidials due to their natural active ingredients, but also have a positive effect on the growth of broiler chickens (Abdul *et al.*, 2009; Abbas *et al.*, 2010; Saleh *et al.*, 2020). *J. mandshurica* has been widely used as a natural medicine to treat various diseases among people for a long time (An-dong *et al.*, 2019; Yuan-Yuan *et al.*, 2019). *J. mandshurica* plant organs are currently known to be enriched with more than 400 compounds, including quinones, phenolics, triterpenoids, diarylheptanoids, flavonoids, coumarins, lignans, phenylpropanoids, steroids, etc (Rami and Hyun, 2006; Yuwei *et al.*, 2016). In addition, various pharmacological activities of *J. mandshurica* were investigated in several studies, and it has been reported that it exhibits various biological activities such as anti-tumor, immunomodulatory, anti-inflammatory, neuroprotective, anti-diabetic, anti-viral, antibacterial and anti-melanin-producing activity (Yao *et al.*, 2012; Yongli *et al.*, 2013; Fei *et al.*, 2021).

Coccidiosis is a well-known protozoan disease in the poultry industry. This disease destroys the intestinal epithelial cells of chickens, which causes bloody diarrhea. When chickens are infected with coccidium, the resistance to the external environment decreases due to a decrease in the immune response, and they display typical clinical symptoms such as loss of appetite and depression (Chapman, 2014; Jabbar *et al.*, 2014). In addition, bloody diarrhea and oocyst leakage through feces are serious problems that can accompany secondary infection and the spread of coccidiosis in poultry (Anteneh *et al.*, 2019). According to the test results, it was found that the aqueous extract of *J. mandshurica* epicarp had anticoccidial properties by inhibiting oocyst activity and inhibiting sporulation. In some studies, the sporulation inhibition rates of leaf and fruit extract of *P. emblica* (1% concentration) were 35.7% and 36.7%, respectively (Usha *et al.*, 2021). This optimum concentration displays the highest inhibition rate

(Usha *et al.*, 2021). However, the sporulation inhibition rates of the aqueous extract of *J. mandshurica* epicarp were 38, 54, 68, 77, and 84% when the concentration were 1 mg/mL (1%), 10 mg/mL (10%), 30mg/mL (30%), 60mg/mL (60%), 100mg/mL (100%), respectively. This revealed that the inhibitory efficiency of the aqueous extract of *J. mandshurica* epicarp against coccidia increases linearly with the concentration increase.

In the broilers fed the aqueous extract of *J. mandshurica* epicarp, the symptoms of bloody diarrhea were significantly lower than that of the PC group, and there was a clear difference in the amount of oocysts excretion through feces. This might be due to the anti-inflammatory and antibacterial effects of the juglone component of *J. mandshurica* epicarp.

Relative body weight gain of experimental groups fed with the aqueous extract of *J. mandshurica* epicarp at 5mL/kg of body weight was higher than that of experimental groups fed the aqueous extract of *J. mandshurica* epicarp at 1, 3, 7mL/kg of body weight. This can be attributed to the various pharmaceutical ingredients of *J. mandshurica* (Yan *et al.*, 2019), which can promote the metabolic activity of the body. As the main medicinal ingredient here, Juglone is a phenolic compound for treating inflammation and infectious diseases. Juglone is active in hepatic metabolism and affects several metabolic pathways related to energy metabolism. Taking a small dose is beneficial to health. However, in some studies conducted on rat liver where Juglone was used in an excessive amount, it was found to interfere with liver energy metabolism by acting as an uncoupler of oxidative phosphorylation in isolated liver mitochondria (Saling *et al.*, 2011). In addition, overdosage or unreasonable use of Juglone can lead to some adverse reactions, such as nausea, vomiting, dizziness, dyspnea, palpitation, and even shock and death (Saling *et al.*, 2011; Huo *et al.*, 2017). This indicates that the aqueous extract of *J. mandshurica* epicarp can promote metabolic activity in vivo at an appropriate dose, but it might also cause some negative effects in vivo when it exceeds a certain amount. Further research in this area should be strengthened in the future. Nevertheless, the relative body weight gain of all experimental groups was significantly higher than the PC group and there were no significant differences in the degree of bloody diarrhea, amount of oocysts excretion through feces, and the relative body weight in the experimental group fed with the aqueous extract of *J. mandshurica* epicarp at 5mL, 7 mL/kg BW compared with the group fed salinomycin. In addition, the results of this study showed that the ACI of groups fed with the aqueous extract of *J. mandshurica* epicarp at 5mL, 7mL/kg of body weight was above 160, which displayed a marked anticoccidial effect.

We have proved that *J. mandshurica* epicarp has a good therapeutic effect on chick coccidiosis, and we

believe that it will become a new anticoccidial medicine because of its low cost, simple pharmaceutical processes, and high application value.

## References

- Abbas RZ, Zafar I, Khan MN, Zafar MA & Zia MA. 2010. Anticoccidial activity of *Curcuma longa* L. in broilers. *Brazilian Archives of Biology and Thchnology*, 53(1):63–67. DOI: 10.1590/S1516-89132010000100008
- Abdul LM, Zhuojian L & Shampa D. 2009. Effect of pine bark (*Pinus radiata*) extracts on sporulation of coccidian oocysts. *Folia Parasitologica*, 56(1): 1-5. DOI: 10.14411/fp.2009.001
- Alaeldein MA, Abdullah HA, Yousif MD & Rifat UK. 2018. The effect of phytochemicals on growth traits, blood biochemical and intestinal histology in broiler chickens exposed to *Clostridium perfringens* challenge. *Journal of Applied Animal Research*, 46 (1): 691–695. DOI: 10.1080/09712119.2017.1383258
- An-dong W, Chao-jie X, Yun-qiang Z, Mei-chen L, Xia W, Jian-yu L & Yong-nan X. 2019.  $\alpha$ -Tetralonyl glucosides from the green walnut husks of *Juglans mandshurica* and their antiproliferative effects. *Planta Medica*, 85: 335–339. DOI: 10.1055/a-0832-2328
- Anteneh W, Ermias M & Yehualashet B. 2019. Prevalence of poultry coccidiosis and associated risk factors in intensive farming system of Gondar Town, Ethiopia. *Veterinary Medicine International*, 2019: 5748690. DOI: 10.1155/2019/5748690
- Applegate TJ, Klose V, Steiner T, Ganner A & Schatzmayr G. 2010. Probiotics and phytochemicals for poultry: Myth or reality? *Journal of Applied Poultry Research*, 19(2): 194-210. DOI: 10.3382/japr.2010-00168
- Arvind SN, Suaib N, Suchita S, Vinay K, Namita G & Mahendra PD. 2011. Antiproliferative and antioxidant activities of *Juglans regia* fruit extracts. *Pharmaceutical Biology*, 49(6): 669-673. DOI: 10.3109/13880209.2010.537666
- Bozkurt M, Selek N, Küçükyılmaz K, Eren H, Güven E, Çatli AU & Çinar M. 2012. Effects of dietary supplementation with a herbal extract on the performance of broilers infected with a mixture of *Eimeria* species. *British Poultry Science*, 53(3): 325-332. DOI: 10.1080/00071668.2012.699673
- Chapman HD. 2014. Milestones in avian coccidiosis research: A review. *Poultry Science*, 93(3):501-11. DOI: 10.3382/ps.2013-03634
- Cobb-Vantress. 2020. Cobb 700 slow feather breeder management supplement (L-013-01-20 EN). Cobb-Vantress, Inc., www.cobb-vantress.com
- Dexing M, Chunli Ma, Long Pan, Guangxing L, Jinghong Y, Jiehua H, Haofan C & Xiaofeng R.

## Acknowledgments

The authors would like to thank the Pyongyang University of Agriculture, DPR of Korea for their assistance in this research.

2011. Vaccination of chickens with DNA vaccine encoding *Eimeria acervulina* 3-1E and chicken IL-15 offers protection against homologous challenge. *Experimental Parasitology*, 127(1): 208-214. DOI: 10.1016/j.exppara.2010.07.015
- Fei L, Ziyang W, Yan Y, Yafei J, Haizhen L, Keqing H, Daoheng L, Xiaofei S, Xirui H & Nan Z. 2021. *Juglans mandshurica* Maxim.: A review of its traditional usages, phytochemical constituents, and pharmacological properties. *Frontiers in Pharmacology*, 11: 1-31. DOI: 10.3389/fphar.2020.569800
- Fernandez S, Pagotto AH, Furtado MM, Katsuyama ÂM, Madeira AMBN & Gruber A. 2003. A multiplex PCR assay for the simultaneous detection and discrimination of the seven *Eimeria* species that infect domestic fowl. *Parasitology*, 127(4): 317-325. DOI: 10.1017/s0031182003003883
- Giannenas I, Florou-Paneri P, Papazahariadou M, Christaki E, Botsoglou NA & Spais AB. 2003. Effect of Dietary oregano essential oil supplementation on performance of broilers challenged with *Eimeria tenella*. *Archives of Animal Nutrition*, 57(2):99-106. DOI: 10.1080/0003942031000107299
- Giannenas I, Tsalie E, Triantafillou E, Hessenberger S, Teichmann K, Mohnl M and Tontis D. 2014. Assessment of probiotics supplementation via feed or water on the growth performance, intestinal morphology and microflora of chickens after experimental infection with *Eimeria acervulina*, *Eimeria maxima* and *Eimeria tenella*. *Avian Pathology*, 43(3): 209–216. DOI: 10.1080/03079457.2014.899430
- Hafez MH. 2008. Poultry coccidiosis: Prevention and control approaches. *Archiv Fur Geflugelkunde*, 72(1) S: 2-7.
- Holdsworth PA, Conway DP, McKenzie ME, Dayton AD, Chapman HD, Mathis GF, Skinner JT, Mundt HC & Williams RB. 2004. World association for the advancement of veterinary parasitology (WAAVP) guidelines for evaluating the efficacy of anticoccidial medicines in chickens and turkeys. *Veterinary Parasitology*, 121(3-4): 189-212. DOI: 10.1016/j.vetpar.2013.02.003
- Huo JH, Du XW, Sun GD, Meng YL & Wang WM. 2017. Comparison of the chemical profiles of fresh-raw and dry-processed *Juglans mandshurica*. *Journal of Separation Science*, 40: 646–662. DOI: 10.1002/jssc.201600877
- Jabbar TA, Chand N, Saddique U, Bailey CA & Khan RU. 2014. Antiparasitic effect of wild rue

- (*Peganum harmala* L.) against experimentally induced coccidiosis in broiler chicks. *Parasitology Research*, 113: 2951-2960. DOI: 10.1007/s00436-014-3957-y
- Jin-Hai H, Xiao-Wei D, Guo-Dong S, Wen-Ting D, Wei-Ming W. 2018. Identification and characterization of major constituents in *Juglans mandshurica* using ultra performance liquid chromatography coupled with time-of-flight mass spectrometry (UPLC-ESI-Q-TOF/MS). *Chinese Journal of Natural Medicines*, 16(7): 525-545. DOI: 10.1016/S1875-5364(18)30089-X
- Johnson J & Reid WM. 1970. Anticoccidial drugs: lesion scoring techniques in battery and floor-pen experiments with chickens. *Experimental Parasitology*, 28(1): 30-36. DOI: 10.1016/0014-4894(70)90063-9
- Kostadinović L, Puvača N, Popovic S & Lević J. 2015. Botanical supplements as anticoccidial alternatives in poultry nutrition. *Worlds Poultry Science Journal*, 71(1): 27-36. DOI: 10.1017/S0043933915000033
- Lan L, Zuo B, Ding H, Huang Y, Chen X & Du A. 2016. Anticoccidial evaluation of a traditional chinese medicine-*Brucea javanica*-in broilers. *Poultry Science*, 95(4): 811-818. DOI: 10.3382/ps/pev441
- Li-yun C, Ke-qian D, Jing X, Yi-fan C, Jian-zhong X, De-He W, Er-ying H, Li-jun X, Hui C & Rong-yan Z. 2021. Effect of natural garlic essential oil on chickens with artificially infected *Eimeria tenella*. *Veterinary Parasitology*, 300: 109614. DOI: 10.1016/j.vetpar.2021.109614
- Mahmood AAQ, Mohamed AD, Esam MAI-S, Mutee M, Mohammed M & Saleh Al-Q. 2020. *Rumex nervosus* leaf extracts enhance the regulation of goblet cells and the inflammatory response during infection of chickens with *Eimeria tenella*. *Journal Of King Saud University Science*, 32(3): 1818-1823. DOI: 10.1016/j.jksus.2020.01.024
- Min Z, Xueyan L, Qiping Z, Rufeng S, Suhan X, Keyu Z, Lifang Z, Xiaoyang W, Mi W, Yingchun L, Chunmei W, Jie Z, Feiqun X & Chenzhong F. 2019. Anticoccidial activity of novel triazine compounds in broiler chickens. *Veterinary Parasitology*, 267: 4-8. DOI: 10.1016/j.vetpar.2019.01.006
- Nahed AE, Mohamed EA, Najah MA, Asmaa FK, Ayman ET, Ayman AS, Mohamed TE, Heba MS, Amira ME, Synan FA, Khaled AE & Ahmed RE. 2022. Phytochemical control of poultry coccidiosis: a review. *Poultry Science*, 101: 101542. DOI: 10.1016/j.psj.2021.101542
- Nilsson O, Greko C, Bengtsson B & Englund S. 2012. Genetic diversity among VRE isolates from Swedish broilers with the coincidental finding of transferrable decreased susceptibility to narasin. *Journal Of Applied Microbiology*, 112(4): 716-722. DOI: 10.1111/j.1365-2672.2012.05254.x
- Peek HW & Landman WJM. 2011. Coccidiosis in poultry: anticoccidial products, vaccines and other prevention strategies. *Veterinary Quarterly*, 31: 143-161. DOI: 10.1080/01652176.2011.605247
- Primož Z & Suzana Ž. 2012. The impact of coccidiostats monensin and lasalocid on Cd and Pb uptake in the isopod *Porcellio scaber*. *Applied Soil Ecology*, 48(1): 36-43. DOI: 10.1016/j.apsoil.2012.01.003
- Qin Y, Qing-Shou Y, Yi K, Yue-Zhi Z, Ling-Ling F, Lu Z, Lin G, Ze-Ping X & Shu-Min Z. 2019. Antimicrobial and cytotoxic juglones from the immature exocarps of *Juglans mandshurica*. *Natural Product Research*, 33(22): 1-7. DOI: 10.1080/14786419.2018.1468326
- Qing L, Ping Z, Xing-Cong L, Melissa RJ, Chong-Ren Y & Ying-Jun Z. 2010. New  $\alpha$ -Tetralone Galloylglucosides from the Fresh Pericarps of *Juglans sigillata*. *Helvetica Chimica Acta*, 93(2): 265-271. DOI: 10.1002/hlca.200900177
- Rami AD & Hyun SL. 2006. Poultry coccidiosis: recent advancements in control measures and vaccine development. *Expert Review of Vaccines*, 5(1): 143-163. DOI: 10.1586/14760584.5.1.143
- Rashed AA & Alaeldein A. 2019. Anticoccidial and antioxidant effects of plants derived polyphenol in broilers exposed to induced coccidiosis. *Environmental Science and Pollution Research*, 26: 14194-14199. DOI: 10.1007/s11356-019-04615-2
- Saleh Al-Q, Mahmood AAQ, Esam MAI-S, Mutee M, Mohammed MM & Mohamed AD. 2020. *Rumex nervosus* changed the oxidative status of chicken caecum infected with *Eimeria tenella*. *Journal of King Saud University Science*, 32(3): 2207-2211. DOI: 10.1016/j.jksus.2020.02.034
- Saling SC, Comar JF, Mito MS, Peralta RM & Bracht A. 2011. Actions of juglone on energy metabolism in the rat liver. *Toxicology and Applied Pharmacology*, 257(3): 319-27. DOI: 10.1016/j.taap.2011.09.004
- Takele B. 2016. Veterinary drug residues in food-animal products: its risk factors and potential effects on public health. *Journal of Veterinary Science and Technology*, 7(01): 1-7. DOI: 10.4172/2157-7579.1000285
- Tuhinadri S & Samir KS. 2015. Medicinal plants, human health and biodiversity: a broad review. *Advances in Biochemical Engineering-Biotechnology*, 147: 59-110. DOI: 10.1007/10\_2014\_273
- Usha NSS, Deepani DF, Kavindra KW, Ariyathilaka M, Indunil P & Rajapakse RPVJ. 2021. Anticoccidial effects of *Phyllanthus emblica* (Indian gooseberry) extracts: Potential for controlling avian coccidiosis. *Veterinary Parasitology Regional Studies and Reports*,

- 25:100592. DOI: 10.1016/j.vprsr.2021.100592
- Wei-Ning B, Wan-Jin L & Da-Yong Z. 2010. Nuclear and chloroplast DNA phylogeography reveal two refuge areas with asymmetrical gene flow in a temperate walnut tree from East Asia. *New Phytologist*, 188: 892-901. DOI: 10.1111/j.1469-8137.2010.03407.x
- Wen-Ting W, Bing X, Da-Yong Z & Wei-Ning B. 2016. Phylogeography of postglacial range expansion in *Juglans mandshurica* (Juglandaceae) reveals no evidence of bottleneck, loss of genetic diversity, or isolation by distance in the leading-edge populations. *Molecular phylogenetics and evolution*, 102:255-264. DOI: 10.1016/j.ympev.2016.06.005
- Xiaohui P, Yao N, Jianjun W, Qiang H & Yuqiang C. 2015. Juglone prevents metabolic endotoxemia-induced hepatitis and neuroinflammation via suppressing TLR4/NF- $\kappa$ B signaling pathway in high-fat diet rats. *Biochemical and Biophysical Research Communications*, 462(3): 245-250. DOI: 10.1016/j.bbrc.2015.04.124
- Yan Z, Wei Z, Shengbao D, Zhe J, Mei J & Gao L. 2019. Phytochemical investigation on the roots of *Juglans mandshurica* and their chemotaxonomic significance. *Biochemical Systematics and Ecology*, 87: 103957. DOI: 10.1016/j.bse.2019.103957
- Yao Y, Yu-Wei Z, Lu-Guo S, Biao L, Yong-Li B, Hua L, Yu Zh, Li-Hua Z, Ying S, Chun-Lei Y, Yin W, Guan-Nan W & Yu-Xin L. 2012. Juglanthraquinone C, a novel natural compound derived from *Juglans mandshurica* Maxim, induces S phase arrest and apoptosis in HepG2 cells. *Apoptosis*, 17(8): 832-41. DOI: 10.1007/s10495-012-0722-5
- Yasuhiro M, Mitsuru K, Noritoshi K & Toshiaki M. 1977. Studies on anticoccidial agents. 10. Synthesis and anticoccidial activity of 5-nitronicotinamide and its analogs. *Journal of Medicinal Chemistry*, 20: 129-133. DOI: 10.1021/jm00211a027
- Yongli Z, Yuqiang C, Jiayong Z, Hongzhi L, Jianwen M, Xiaobao J, Xiangsheng W, Yifan D & Jiazheng L. 2013. The anti-tumor effect and biological activities of the extract JMM6 from the stem-barks of the Chinese *Juglans mandshurica* Maxim on human hepatoma cell line BEL-7402. *African Journal of Traditional Complementary and Alternative Medicines*, 10(2): 258-269. DOI: 10.4314/ajtcam.v10i2.10
- Yuanyuan Z, Bingyou Y, Zhaoxi L, Yanqiu J, Yuxin L, Lei F, Xiaoli W & Haixue K. 2015. Cytotoxicity of Triterpenes from Green Walnut Husks of *Juglans mandshurica* Maxim in HepG-2 Cancer Cells. *Molecules*, 20(10): 19252-62. DOI: 10.3390/molecules201019252
- Yuan-Yuan Z, Shuang G, Ying W, Hong-Juan S, Hui-Rui G, Xiao-Juan Z, Yan-Ping S, Yan L, Bing-You, Y & Hai-Xue K. 2019.  $\alpha$ -Tetralone glycosides from the green walnut husks of *Juglans mandshurica* Maxim. and their cytotoxic activities. *Natural Product Research*, 34(13): 1-9. DOI: 10.1080/14786419.2018.1561681
- Yuwei Z, Hua L, Shanshan L, Jianbo C, Yinshi S & Yuxin L. 2016. High-speed counter-current chromatography assisted preparative isolation of bioactive compounds from stem bark of *Juglans mandshurica*. *Journal of Separation Science*, 40(3): 767-778. DOI: 10.1002/jssc.201601043