

Comparison of Measurement Indices of Diversity, Richness, Dominance, and Evenness in Rangeland Ecosystem (Case Study: Jvaaherdeh-Ramesar)

Negar Davari^A, Mohammad Hassan Jouri^B, Ali Ariapour^C

^AMSc. Student of Range management, Islamic Azad University, Noor branch, Email: davarnegar63@yahoo.com

^BAssistant Prof. Dept. of Natural Resources, Islamic Azad University, Noor Branch. Iran.

^CAssistant Prof. Dept. of Natural Resources, Islamic Azad University, Boroujerd Branch. Iran.

Manuscript Received: 30/09/2010

Manuscript Accepted: 11/11/2011

Abstract. The diversity and richness of species are of important components to evaluate the health of ecologic system of rangelands. This research was conducted to investigate the biodiversity in riparian and neighboring areas via comparison of indices in Jvaaherdeh rangeland of Ramsar. Plot-transects was randomly used to estimate diversity, richness, dominance, and evenness indices in two regions. Multi Linear regression method was used to analyze the relationships of three indices to rangeland condition. Results of floristic list showed that there were 66 species belong to 19 families and 56 genera. Gramineae and Compositae families with values of (23.24 and 12.1%) had the highest and Boraginaceae and Hypolepidaceae with 2% had the lowest frequency, respectively. Result showed that richness species index in riparian area and diversity index in neighboring area had high correlation with the rangeland conditions. Hence, ecologic management could be used as a tool for evaluation of indices as reflectors of disturbance in rangeland.

Key words: Indices, Diversity, Richness, Dominance, Evenness, Riparian area, Mountainous rangeland, Regression.

Introduction

An ecologist has to have correct knowledge about the function of different ecosystems by which he/she could manage them ecologically. There are different methods to investigate ecosystem components. Knowledge about natures and functions of ecosystem could help one to having a suitable ecosystem management (Jacobs, 1975). In most of the published studies, species variation as a biological diversity (DeLong, 1996; Williams, 2004) consists of two components: species richness and evenness (Liu *et al.*, 2008, and Omernik, 2003) which can be regarded as sustainability and healthiness factors in the rangeland ecosystems (Williams, 2004; Jouri, 2010). Different factors have been already affected by the community's diversity as follows: sampling size and rate; destruction and human activity, farming, topographical factors, succession, underground waters, rivers and livestock grazing (Afshani *et al.*, 2009). The riparian areas, which held the highest amount of the above situations in rangeland level, can be considered as one of the most sensitive areas where need an ecological and sustained management (Khatibi, 2005, Williams, 2004). Whereas the amplex concept and situation as well as the species variety have not been understood correctly (Hooper and Vitousek, 1997). Therefore, it needs a wide range of surveys to show the relationships among the mentioned components in an ecosystem. Through this, the researchers can investigate plant species diversity to identify their status by considering management role in different plant colonies (Goodman, 1975; Maguran, 1996). Evaluation of diversity parameters like species combination, dominance, evenness and the amount of species can be used for estimating the ecological status (Goodman, 1975). Identifying ecosystem health, Encoist *et al.* (2002) have pointed out that there are sorts of positive relation among the

number of genus, family and species in all sampled units (Enquist *et al.*, 2002) in such a way, increasing of species richness would bring more diversity in genus and family. The results gained by Lorio (2002) showed that livestock grazing, livestock intensity and grazing systems could affect plant diversity so that knowledge of diversity changes has an important role to determine the rangeland conditions of the rangeland ecosystem and managing the impacts. Wilcox *et al.* (1987) studied how different kinds of grazing intensities can affect plant communities. Moreover, they concluded that if there would be moderate grazing in contrast to light and over grazing, it never leads to an increase in the species richness. Light grazing up to mid level could increase the species diversity and also homogeneity in rangeland plant combinations, however, overgrazing would decrease the important rangeland species and if there won't be happened any grazing, the dominant species could occupy all over the area. It has been expressed that there is a kind of direct relationship between species diversity and richness to the rangelands conditions. By surveying of diversity in the rangeland ecosystems, one will be able to know about the rangeland conditions and then use it as an index in management and programming of rangelands (Gillison, 2006; Ludwig & Renolds 1988; Menhinick, 1964; Pimm, 1984). Regarding this, Mahdavi *et al.* (2005) in investigation of the rangeland conditions in Inchebron region using diversity and richness indices, have been found out that the Shannon diversity's index and richness index of Margalof could show a better description for the conditions of rangeland health. With reference to the rangeland analysis, riparian areas as a part of the rangeland ecosystems, had varieties of species which are mesic plants. Secondly, it has high diversity because of its role as buffer zone between water and ground areas in such a way that

the biological activities at this ecosystem would be so dynamic (Brodt *et al.*, 2009). Some of managing and human activities on the rangeland ecosystems can lead to decreasing and demolishing of natural habitats and resilience of species diversity and richness, therefore, it require a strong managing strategy in a long time (Hawksworth, 1995). However, vegetarian evenness in nature could be easily reversible (Orians, 1975; Shaojun *et al.*, 2001 and Smith, 2001). There is a direct relation between species diversity and richness in mountainous areas (Jouri *et al.*, 1381.) and their increase also could bring more stability, species diversity and raise the forage quality of range species (Pielou, 1975). Consequently, comparing the biological diversity indices and their

efficiency to introduce the subjective facts of the rangeland ecosystems can be an important step in preparing and introducing different managing approaches that it is the aim of this research.

Materials and Methods

In order to compare the biological diversity indicators in riparian ecosystems, the rangeland areas in Javaherdeh (Ramsar) that its geographical coordinates are 36°54'N and 50°40'E in northern Alborz was selected. The climate of the area varied from cold and humid condition (in lower altitudes) up to altitude climate in higher areas with annual average of rainfall recorded as 640 mm (Jouri, 2010).




Fig. 1. Position of the study area in the Province and town

Research Approach

First of all, using a topographical map in scale of 1:25000 and field monitoring, the territory of study area was determined and the landform units were derived. To evaluate the function of riparian area safety and to consider the V shape of the area, we prepare a cross section of the

area and the neighborhood region to determine the diversity and species richness (Schade *et al.*, 2001). Then, a floristic list was prepared and completed through collecting and identifying them by color Flora and Flora of Iranica (Ghahraman, 2000).

Table 1. Names of some of species in the area investigated coming through floristic list

Number	Species name	Family	Biological type
1	<i>Achillea millefolium</i>	Compositae	Ch
2	<i>Cardamine pratensis</i>	Cruciferae	Tr
3	<i>Carex ovalis</i>	Cyperaceae	-
4	<i>Agropyron intermedium</i>	Gramineae	Ch
5	<i>Chenopodium album</i>	Chenopodiaceae	Tr
6	<i>Alchemilla persica</i>	Rosaceae	Ch
7	<i>Dactylis glomerata</i>	Gramineae	Ch
8	<i>Echinops robustus</i>	Compositae	Ch
9	<i>Alopecurus sp</i>	Gramineae	Ch
10	<i>Euphorbia helioscopia</i>	Euphorbiaceae	Ch
11	<i>Anchusa iranica</i>	Boraginaceae	Ch
12	<i>Anthemis cotula</i>	Compositae	Ch
13	<i>Arctium lappa</i>	Compositae	Ch
14	<i>Artemisia aucheri</i>	Compositae	Ch
15	<i>Plantago major</i>	Plantaginaceae	Cr
16	<i>Astragalus sp</i>	Fabaceae	Ch
17	<i>Avena barbata</i>	Gramineae	Tr
18	<i>Malva neglecta</i>	Malvaceae	Tr
19	<i>Onobrychis cornuta</i>	Fabaceae	Ch
20	<i>Poa annua</i>	Gramineae	Tr
21	<i>Poa bulbosa</i>	Gramineae	Cr
22	<i>Bromus persica</i>	Gramineae	Ch
23	<i>Bromus tectorum</i>	Gramineae	Tr
24	<i>Poterium sanguisorba</i>	Rosaceae	Ch
25	<i>Capsella bursa-pastoris</i>	Cruciferae	Tr

The sampling method in riparian and adjacent areas has been done randomly (Body & Svejcar, 2004; Clary, 1995) using transect-plot method (Robins *et al.*, 2001; Jouri, 2010; 73, Sabetpour *et al.*, 2002) and the determination of plot size was done by Minimum Area method (Cain, 1932) and the plot number also was obtained by statistical method (Pielou, 1975). Three different transects from 20 to 50 meters were used in the study area (Robins *et al.*, 2001). The plots must be randomly settled along with the big diameter of the investigated area (Smith and Wilson, 1996). Sampling was done in the mid of flowering period in each

stand area individually. The vegetation traits, frequency and the amount of species were recorded for each plot to infer the biological diversities' indices (Table 2). Calculating the biological indicators was done through PAST v.2.7 software. In order to find out the relationship between biological indices and the rangeland conditions, multivariate statistics was used via stepwise regression method using SPSS v.17 software (Afshani *et al.*, 2009). In this research, the rangeland conditions and biological indicators were recognized as dependent and independent variables, respectively.

Table 2. Formula of diversity, richness, and evenness and dominance indexes (adapted of Ejtehadi et al., 2009)

The Changes range	Formula	Component	Index
0-1	$H' = -\sum_{i=1}^s p_i \ln p_i$	Shannon-Wiener	
0-1	$D = \sum_{i=1}^s P_i^s$	Simpson	
0-1	$d = N_{max} / N$	Berger-Parker	
0-1	$(ax), \frac{(ax)^r}{r}, \frac{(ax)^r}{r}, \dots, \frac{(ax)^n}{n}, \dots$	Fisher alpha	
0 - ∞	$D_{mg} = \frac{S-1}{\ln N}$	Margalef	
0 - ∞	$D_{Mn} = \frac{S}{\sqrt{N}}$	Menhinick	
0-1	$\hat{D}_{Max} = \frac{1}{s}$	Simpson	Evenness
0-1	$C = \sum (ni/N)^2$	Simpson	Dominance

Results

According to the gained results, 66 species were recognized that all were belong to 19 families and 56 genera. The families of Gramineae and Compositae had the most frequency (12.1% & 23.2%) and the families of Boraginaceae and Hypolepidaceae with 2% compose the species coverage of the study area (Fig.

2). Therefore, in Gramineae family, *Bromus tomentellus* shows the most abundance (18.75%) and *Digitaria* sp. with the least amount (1%) as the area flora. The results of classifying biological type in Raunkiaer method in the area showed that Hemicryptophyte (70.73%), Therophyte (20.33%), Geophyte (8.13%) plants had the most amounts of biological forms in the area (Fig. 3).




Fig. 2. Vegetarian families in Javaherdeh




Fig. 3. The biological type in Javaherdeh

In order to signify the exploring of biological diversity indices in both riparian and adjacent areas, T-test was used. The result showed that the richness index of Margalef was the only significant index in the both areas (Table 4).

Furthermore, in each region, the

biological indices were individually analyzed along with rangeland conditions. The results of multivariate regression analysis in riparian area showed that both Margalef and Menhinick indices (from species richness class) had significant relationship with the rangeland conditions (Table 5).

Table 4. The amount of bio-diversity indices in both areas: riparian and adjacent

Zone	Average \pm (standard division)	T statistic	Component	Index
Riparian	2.095 \pm 0.356	0.767	Shannon-Wiener	Diversity
Adjacent	2.174 \pm 0.389	0.766		
Riparian	0.837 \pm 0.076	-1.348	Simpson	
Adjacent	0.847 \pm 0.084	-1.346		
Riparian	2.839 \pm 0.862	-0.767	Berger-Parker	
Adjacent	3.161 \pm 1.159	-0.766		
Riparian	0.255 \pm 0.113	-1.987	Fisher-alpha	
Adjacent	0.245 \pm 0.118	-1.982		
Riparian	2.21 \pm 0.577	-1.820 *	Margalef	Richness
Adjacent	2.23 \pm 0.706	-1.815 *		
Riparian	0.870 \pm 0.206	-0.602	Menhinick	
Adjacent	0.942 \pm 0.258	-0.601		
Riparian	0.866 \pm 0.869	-2.038	Simpson	Evenness
Adjacent	0.875 \pm 0.098	-2.031		
Riparian	0.162 \pm 0.076	0.568	Simpson	Dominance
Adjacent	0.153 \pm 0.084	0.568		

* = significant at 5% level

Table 5. Multivariate regression in riparian area on basis of stepwise method

Index	Coefficient of Correlation R	Coefficient of determination R ²	F-statistic	Sig.
Margalef	0.252	0.064	5.584	0.000
Menhinick	0.428	0.182	9.080	0.001

The regression equation (1) of both variables considering the rangeland conditions is as follows:

$$Y=60.973-59.052X_1+26.969X_2(1)$$

In addition, the results show that the amount of Margalef index in defining rangeland condition's score changes is more observable than the others (regression model 2).

$$Y=8.19X_1-0.66X_2$$

The above equation shows that the most changes are related to Margalef richness (positive), and then it follows the Menhinick one (negative).

The regression results out of river adjacent area show that the studied indices of Shannon and Fisher-alpha (diversity indices) and Menhinick richness in contrast to the others show sorts of significant relation to rangeland score (Table 6).

Table 6. Multi variable regression in river adjacent area

Index	Coefficient of Correlation R	Coefficient of determination R ²	F-statistic	Sig
Shannon	0.414	0.171	5.265	0.001
Fisher alpha	0.547	0.200	11.127	0.000
Menhinick	0.550	0.202	8.222	0.000

While the regression equation of this analysis was as follows:

$$Y=2.118-0.145X_1+0.16X_2-0.55X_3$$

The standard model shows that the amount of Fisher-alpha index (directly & positively) and Menhinick richness (inverse) had the most and Shannon index (inverse) the least role in defining the rangeland changes (below regression model):

$$Y=-0.54X_1-1.38X_2+1.84X_3$$

Discussion and Conclusion

Reserving the biodiversity as a natural heritage can be challenged mostly today by politicians and scientist as well as preserving the flora species diversity as the only productive creatures besides being the bio-bed for the other organisms can be regarded as the most important ones in contrast to the other biodiversity components. To calm all these debates, we must analyze the bio indices. Regarding that, the area flora in altitude of 1800 to 3000 m was remarkable (Fig. 2), but comparing the bio-indices analysis

shows that the Menhinick and Margalef indicators had the most correlation to the rangeland condition's score. Having a high degree of just one genus species (*Bromus*) in an area shows that the utilize factors (livestock & human) had interfered somehow in unifying the area flora. The field observations note that riparian area had been severely affected by free livestock grazing such as sheep, goat, cow or sometimes horses. Furthermore, it is not so overstated that the wetlands such as the riparian area in possession of wet and succulent species can attract many livestock in contrast to the other areas. Therefore, the existence of palatable and semi-palatable species would be severely affected by different kinds of livestock and animals in the area because of their different feed requirements so that according to the observation, these species can be found refuge in spiny shrubs among the rocks and the steep slopes. So, it is not an imagination that species richness in both areas could be related mostly to rangeland conditions more than the other

bio-indices which for sure can be along with the gained results by West (1993) and Lorio *et al.* (2002). Furthermore, through analyzing bio-indices in the adjacent areas (off the water margin), we have got that the diversity indicators have already shown the most amount of changes in the rangeland conditions. Being away from the riverbank and water resources causes less appearance of livestock at these areas, so lots of different flora species are seen here. Diversity species index therefore can be more significant than the others can (Table 6). In addition, in some steps when the animals are mostly attended, the species richness can be accounted for different changes in status that this part according to field monitoring is clearly obvious. Then, the gained results are completely in accordance with the results coming out of Robines *et al.* (2001) and Jouri (2010). Considering the results coming of analyzing, the rangeland conditions and studying bio-indices draw an acceptable view in account for the health of rangeland ecosystem that can be used in defining the ecologic management of the rangeland. Therefore, using bio-indices can be regarded as a suitable ecological indicator in analyzing the rangeland ecosystems that is in agreement with the results in Odum (1969), Jouri *et al.* (2008) and Mahdavi *et al.* (2005) studies.

References

- Afshani, S. A., M. Norani, M. Mahdavi & A. Norian, 2009. Applicable Authority SPSS₁₇, Bishe Application, 437pp. (In Persian).
- Baghani, M., A. Spehri and F. Fadaei, 1388. Taxonomic surface of plant as a determining factor in diversity community. (Case study the rangeland of pilgrim area in Gorgan, Golestan province) *Jour. Iran desert and rang.* 98 pp. (In Persian).
- Bihamta, M. R. & M. A., Zare Chahoki, 2010. Statistics principles in natural resources. Tehran University Publication. 300 pp. (In Persian).
- Body, C. S. & T. J. Svejcar, 2004. Regrowth and production of herbaceous riparian vegetation following defoliation. *Range management*, **57(5)**: 448-454.
- Brodts, S., K. Klonsky., L. Jackson & S.B. Brush, 2009. Factors affecting adoption of hedgerows and other biodiversity enhancing features on farms in California, USA. *Agrofor Syst*, **76**: 195-206.
- Cain, S. A., 1932. Concerning certain ecological concepts. *Ecological Monoger*, **27(2)**: 475-508.
- Clary, W. P, 1995. Vegetation and soil responses to grazing simulation on riparian meadows. *Range Management*, **48(1)**: 18-25.
- DeLong, D. C. 1996. Defining biodiversity. *Wildlife Society Bulletin* **24**:738-749.
- Ejtehad, H., A. Sepehri & H. R. Akefi, 2009. The Measuring way of biodiversity. Mashhad Ferdosi Publications University. 229pp. (In Persian).
- Enquist, B. J., J. P. Haskell & B. H. Tiffney, 2002. General patterns of taxonomic and biomass partitioning in extant and fossil plant communities. *Nature*, **419**: 610-613.
- Ghahraman, A, 2000. Colored flora of Iran. Vol. 1-22. Forest and Rangeland. (In Persian).
- Gillison, A. N., 2006. A field manual for rapid vegetation classification and survey for general purposes. *International Forestry*, 3-10.
- Goodman. D. 1975. The theory of diversity-stability relationships in Ecology. *The Quarterly Review of Biology*, **50**: 237-266.
- Hawksworth D. L, 1995. Biodiversity:

- Measurement and Estimation. Chapman and Hall, London.
- Hooper, D. U & P. M. Vitousek, 1997. The effects of plant composition and biodiversity on ecosystem processes. *Science*, **277**: 1302–1305.
- Jacobs, J, 1975. Diversity Stability and maturity in ecosystems influenced by human activities. In: van Dobben, W. H. & Lowe-McConnell, R.H. (eds.) *Unifying Concepts in Ecology*. Junk, The Hague, 187-207.
- Jouri, M. H., B. Tomzad. M. Shokri & Banihashemian, 2002. Comparison between diversity and riches indicators in health evaluating of range land. *Iranian jour. Rangeland*. **2(4)**: 344-356. (In Persian).
- Jouri, M. H, 2010. Ecological investigation of upland rangelands (Alborz Mountain) in scale of two Phytogeographical regions of Irano-Touranian and Euro-Siberian, Pune University, India, 960p. (In Persian).
- Khatibi, R., 2005. Ecologic view on range analysis of Taftan, case study: Dejing range of Khash, MSc. Thesis, Mazandaran University, Natural Resources faculty, 106 PP. (In Persian).
- Liu, Z. F., GH. Liu., B.J. Fu & X.X. Zheng, 2008. Relationship between plant species diversity and soil microbial functional diversity along a longitudinal gradient in temperate grasslands of Hulunbeir, Inner Mongolia, *China. Ecol. Res.*, **23**: 511–518.
- Loreau, M., S. Naeem, & P. Inchausti, 2002. Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science*, **294**: 804-808.
- Ludwig, J. A. & Renollds J. F. 1988. *Statistical Ecology: A Primer on, Methods and computing*. John Wiley & Sons.
- Lyon, J. & C. L. Sagers, 2002. Correspondence analysis of functional groups in a riparian landscape. *Plant Ecol*. **164**: 171-183.
- Maguran, A. E, 1996. *Ecological diversity and its measurement*. Chapman & Hall.
- Mahdavi, M., M. R. Chaeichi & M. H. Jouri, 2005. The comparison of evaluating rangeland way selects the better way in North east Iran. (Case study: Golestan Inchebron Rangeland). *Agriculture Science Research And Khazar Natural Resources*, **2(3)**: 42-45 (In Persian).
- Margalef, R, 1957. Diversidad de especies en las commundades naturales. *publications del instituto de biological aplicate*. **6**: 59-72.
- Menhinick, E. F. 1964. A comparison of some species individual diversity indices applied to samples of field insects. *Ecology*, **45**: 839-861.
- Odum, E. P., 1969. The strategy of ecosystem development, *Science* **164**: 262–270.
- Omernik, J. M., 2003. The misuse of hydrologic unit maps for extrapolation, reporting, and ecosystem management. *Jour. Am. Water Resources Assn*. **39(3)**: 563-573.
- Orians, G. H. 1975. Diversity, stability and maturity in natural ecosystems. In: van Dobben, W. H. & Lowe-McConnell, R. H. (eds.) *Unifying Concepts in Ecology*. Junk, The Hague, 139-50.
- Pielou, E. C., 1975. *Ecological Diversity*. Wiley, New York.
- Pimm, S.L, 1984. The complexity and stability of ecosystems. *Nature*, **307**: 321-326.
- Robins P., R. B. Holmes & K. Laddish, 2001. *Bringing farm edges back to life*. Yolo County Resource Conservation District, Woodland, CA.

- Sabetpour, T., M. H. Jouri & R. Erfanzadeh, 2002. Preliminary study to selection of the best method of range condition's determination on upland rangeland of north of Iran, proceeding of third conference, technical Publication of rangeland and range management, **369**: 930-946. (In Persian).
- Schade, J. D., S. G. Fisher, N.B. Grimm & J.A. Seddon, 2001. The influence of a riparian shrub on nitrogen cycling in a sonoran desert stream. *Ecological Society of America*, **82(12)**: 3363-3376.
- Shannon, C. E., 1948. A mathematical theory of communication. *Bell system Technical Jour*, **27**: 379-423.
- Shaojun, X., N. C. Hrister & J. Matse, 2001. Effects of litter accumulation on riparian vegetation: importance of particle size. *Vegetation Science*, **12(2)**: 231-236.
- Simpson, E. H. 1949. Measurement of diversity. *Nature*, **163**: 688 p.
- Smith B. and Wilson J. B., 1996. A Consumer's guide to evenness indices. *Oikos* **76**: 70-82.
- Smith, S. J. 2001. Rethinking riparian regrowth. *Rangelands*, **23(3)**: 1-4.
- West, N. E., 1993. Biodiversity of rangelands. *Jour. Range Management* **46**: 2-13.
- Wilcox, B. P., F. C. Bryant & B. Fraga, 1987. An evaluation of range condition on one range site in the Andes of central Peru. *Jour. Range Management*, **40**: 41-45.
- Williams, J., 2004. Metrics for assessing the biodiversity values of agricultural landscapes. *Pacific Conservation Biology*, **10**: 145-163.