

Contents lists available at Sjournals



Journal homepage: www.Sjournals.com



Original article

Histomorphometric study of the prenatal development of the kidney of one-humped camel (*Camelus dromedarius*) in north-western Nigeria

A. Bello*^a, U.F. Lawal^a, M.L. Sonfada^a, M.A. Umaru^b, J.E. Onu^a, S.A. Shehu^a, A.A. Umar^a

^aDepartment of Veterinary Anatomy, Usmanu Danfodiyo University, Sokoto, Nigeria.

^bDepartment of Theriogenology and Animal production, Usmanu Danfodiyo University, Sokoto, Nigeria.

*Corresponding author; Department of Veterinary Anatomy, Usmanu Danfodiyo University, Sokoto, Nigeria, Tel: +234(0)8039687589.

ARTICLE INFO

Article history:

Received 12 April 2013

Accepted 20 April 2013

Available online 25 April 2013

Keywords:

Histomorphometry

Camel

Kidney

Prenatal development

Northern Nigeria

ABSTRACT

This study aimed at investigating the prenatal development of the kidney of the camel using standard histomorphometric methods. In the experiment, fifteen foetuses obtained from Sokoto metropolitan abattoir at different gestational ages were used for the study. The fetuses were weighed and grouped and aged. In all the foetuses the kidneys were bean-shaped with smooth external surface and thick fibrous capsule which becomes thicker with advancement of gestational age. Well developed ureters and distinct renal pelvis were evident at all stages of development. In each gestational age, values of the weight, length, width and volume of the kidney were determined. In addition the periodic differentiation of the cortex and medulla; their significance in relation to the renal histodifferentiation was discussed. Histological sections showed evidence of collecting duct system that are numerous and extensively coiled in the cortex and medulla of the third trimester kidneys. This is an indication of the ability of the kidney to concentrate urine, in accordance with the counter-current mechanism of the kidney collecting system. It was suggested that the kidney's of the camel possesses the anatomical features for the production of hypertonic urine owing with the adaptive behavior of desert animals. Thick capsule that increases in thickness was also noticed in the kidneys.

1. Introduction

The one-humped Dromedary (*Camelus dromedarius*) and the two-humped Bactrian (*Camelus bactrianus*) camels are among the largest mammalian species, adapted to the desert with its high temperatures and extreme desiccation (Bello et al., 2012). Camelids have been known to be capable of withstanding harsh conditions characteristic of the semi-arid and arid regions because of their peculiar physiologic and morphologic features (Bello et al., 2008). Development of sophisticated physiological adaptations for coping with high temperature, feed and water scarcity in the dry and rough habitat of the camel are seen. These adaptations seem to depend on its ability to tolerate severe dehydration and to economise the meagre water availability (Aly 2007). During the last century, specific investigations have highlighted the mechanisms governing such adaptation including anatomical and physiological peculiarities (Aly 2007).

The kidney is an important organ involved in the removal of unwanted nitrogenous substances, excess water and relative maintenance of osmotic concentration of the blood (Salehi and Morovati, 2012). Sperber (1994) was one of the first to note that certain features involving renal anatomy in different mammals vary with the aridity of the habitat. Numerous textbooks of anatomy and histology (Weibel, 1963a; Wirz. and Dirix, 1973; Fawcett and Raviola, 1994; Eurell et al., 2006; Dyce et al., 1995; Sisson, 1975) have described the anatomy and histology of the adult kidney of domestic animals, and also research studies that investigated kidney development and morphology. The morphometric observations on the kidney of camel (*Camelus dromedarius*) (Abdalla & Abdalla, 1979), one horned rhinoceros (Talukdar et al., 2003), biometrical study of the kidney of buffalo (Malik et al., 1978), morphometric study on kidney of African rat fetus (Onyeanusu et al., 2006) and measurement of renal dimension (Moorthy and Venugopal, 2001). With the changing scenario of camel management, investigations into the adaptation of camel is important (Kataria et al., 2007), with good emphasis on the prenatal development of the kidney is considerable.

Most available reports on renal anatomy were entirely concerned with adult structures of the kidney, thus there is paucity of information about the prenatal development of camel kidney in this area. In this research, gross changes, morphometric changes and histological differentiation involved in the development of the kidney in one-humped camel will be described. The study will also add to the existing information on the morphometric analysis in camels. Therefore this investigation was carried out to reveal and evaluate the pattern of the kidney growth during prenatal life in camel (*Camelus dromedarius*) fetuses.

2. Materials and methods

The study was carried out on 15 fetuses of the one-humped camel collected from the metropolitan abattoir, Sokoto at different gestational ages. The collected fetuses were then taken to the Veterinary Anatomy laboratory of Usmanu Danfodiyo University; where the weight and age of the foetus were determined. The foetal body weight was measured using electrical (digital) weighing balance for the smaller fetuses and compression spring balance (AT-1422), size C-1, sensitivity of 20kg X 50g in Kilogram for the bigger fetuses. The approximate age of the fetuses was estimated by using the formula adopted by El-wishy et al., (1981) using Crown Vertebral Rump Length (CVRL). Based on this, samples were divided into 3 main groups as

1st trimester = below 130 days

2nd trimester = 131- 260 days

3rd trimester = 261 - 390 days

The fetuses were then placed on a dorsal recumbency and a mid-line incision was made, starting from the xiphoid cartilage and extending to the pubic symphysis. The peritoneum was reflected and the intestine displaced to gain access to the reproductive organs. The organs were examined in situ and exteriorised. The length, weight,

and width of each kidney were measured using ruler and thread, weighing balance, and vernier callipers, respectively. The length was taken from the cranial pole to the caudal pole along the longitudinal axis while the width was taken from the median fossa (hilus) to the lateral edge of the hilus. The Volume of each kidney was determined by water displacement technique (Archimedes principle). The organs were weighed using an electronic weighing (Mettler® balance P1210, Mettler Instruments AG, Switzerland) with a sensitivity of 0.01 g. The data obtained were expressed as mean \pm standard error of the mean (mean \pm SEM).

1cm² thick of sample from each the medulla and cortex was collected and fixed in 10% formalin solution. After fixation was achieved, the tissue sample was processed for paraffin blocks preparation. The sections of 5 μ m were subjected to haematoxylin and eosin for routine morphology (Luiz and Jose, 2005). The standard sections were examined under light microscope and micrographs taken using Sony digital camera(x5) with 12.1 mega pixel.

3. Results

A total of fifteen camel foetuses were used and grouped into first, second and third trimesters comprising of five, seven and three foetuses respectively as shown in Table 1. In all stages of development, the kidneys were found to be irregularly elongated and reddish-brown in colour. Both kidneys were bean-shaped, having a smooth surface and covered by a thin fibro-muscular capsule (which tend to become whiter with the advancement of gestational age) as shown in Plate III. The kidneys were located below the transverse processes of lumbar vertebrae on each side, with the right kidney being more cranial in all camel foetuses as shown in plate II. At the hilus of each kidney adipose tissue were found. Biometrical parameters were shown to be increasing with advancement of gestational age as shown in Table 2. The mean weight, mean length, mean width and mean volume of the right and left kidneys were determined as shown in Table 2.

Table 1

The Mean CRVL, Mean Weight and Sex of the camel foetuses at various trimesters of age.

Trimester	Mean CRVL (cm \pm SEM)	Sex		Mean weight (g \pm SEM)
		Male	Female	
1 st	15.30 \pm 1.74	3	2	68 \pm 21.49
2 nd	40.42 \pm 5.35	3	4	1990 \pm 733.17
3 rd	93.00 \pm 1.52	2	1	11600 \pm 378.59
Total		8	7	

Table 2

The Mean \pm SEM of biometrical parameters of right and left kidneys in relation to stages of development.

Age of the foetus	1 st trimester	2 nd Trimester	3 rd trimester
Sample size	5	7	3
Mean weight(g \pm SEM)			
Right Kidney	0.48 \pm 0.19	8.44 \pm 4.56	49.13 \pm 2.11
Left Kidney	0.48 \pm 0.19	8.71 \pm 4.55	50.33 \pm 1.42
Mean length(cm \pm SEM)			
Right Kidney	1.34 \pm 0.16	3.11 \pm 0.58	6.90 \pm 0.10
Left Kidney	1.32 \pm 0.12	3.04 \pm 0.55	7.06 \pm 0.08
Mean width (cm \pm SEM)			
Right Kidney	0.80 \pm 0.08	1.77 \pm 0.30	3.53 \pm 0.12
Left Kidney	0.78 \pm 0.09	1.7 \pm 0.26	3.70 \pm 0.05
Mean volume (ml \pm SEM)			
Right Kidney	0.40 \pm 0.07	4.21 \pm 1.27	12.4 \pm 0.31
Left Kidney	0.41 \pm 0.09	4.39 \pm 1.27	12.6 \pm 0.31

Histological observation shows that the camel kidney appeared to have the two zones of cortex and medulla right from 1st trimester as shown in Figures 1-6. The development of renal tissues were found to be in succession, from the stage of mesenchyme cells, developing glomerulus and finally to fully developed glomerulus with advancement of gestational age among the three groups as shown in figures 1-6. The duct system also followed the same trend of development from immature to matured duct system. The cortex at first trimester revealed numerous mesenchymal cells and connective tissue with few developing glomerulus (Fig. 1). At second trimester few fully developed glomerulus were found, moderate number of mesenchyme cells and immature developing duct system (Fig. 2). Numerous developed glomerulus and matured duct system with moderate developing glomerulus and connective tissue were seen at third trimester. There was extensive coiling of the duct system at the cortex of the kidney (Fig. 3). The medulla in the camel kidney at first trimester shows only various types of connective tissue (collagen, reticular and elastic) fibers and mesenchyme cells (Fig. 4). At second trimester the medulla was found to have numerous developing collecting ducts, few developed collecting ducts and few mesenchyme cells (Fig. 5). Developed collecting ducts were predominant in the medulla at third trimester with few mesenchyme cells (Fig. 6).



A



B

Plate A. Dissected first trimester camel foetus showing the location of foetal kidneys in situ (Black arrows).

Plate B. Camel foetal kidneys showing smooth external appearance and bean-shaped right kidney (R) and left kidney (L).




Fig. 1. Photomicrograph of Camel kidney (cortex) at 1st Trimester showing developing glomerulus (pG), connective tissue (C), blood vessels (Bv), tunica adventitia (Ta) and mesenchyme cells (Mc) H&E x200.




Fig. 2. Photomicrograph of Camel kidney (cortex) at 2nd Trimester showing glomerulus (G), connective tissue (cT), developing collecting ducts (Pd), kidney capsule (C) and mesenchyme cells (Mc) H&E x200.




Fig. 3. Photomicrograph of Camel kidney (cortex) at 3rd Trimester showing developed glomerulus (G), developing glomerulus (dG), connective tissue (cT) and developing collecting ducts (Cd). H&E x 200.




Fig. 4. Photomicrograph of Camel kidney (medulla) at 1st Trimester showing connective tissues collagen (green arrow), reticular (black arrow), elastic (blue arrow) and mesenchyme cells (yellow arrows) H&E x200.




Fig. 5. Photomicrograph of Camel kidney (medulla) at 2nd Trimester showing developing collecting ducts (Cd), developed collecting ducts (C) and mesenchyme cells (Mc) H&E x200.




Fig. 6. Photomicrograph of Camel (medulla) at 3rd Trimester showing developed collecting ducts (mC) and mesenchyme cells (M) H&E x200.

4. Conclusion

From the study, the result shows that with the advancement of gestation the morphometric data were increasing progressively. This is in accordance with the finding of Hena et al., 2012, on pigeon and Bello et al., 2012 on camel digestive tract.

All kidneys were found to be irregularly elongated and reddish-brown in colour. Both kidneys are bean-shaped, having a smooth surface and covered by a thin fibro-muscular capsule. This is in accordance with the reports of Smuts and Bezuindenhout, (1987); Dyce, (1995); Abdalla et al., (1974); Dowelmadina, (2012); Salehi et al., (2012).

Both kidneys were located below the transverse processes of lumbar vertebrae of each side, with the right one being more cranial in all groups. Abdalla et al., (1974) reported that right kidney is situated under the first three lumbar vertebrae while the left one is found under the last three lumbar vertebrae. Salehi et al., (2012) also reported that the right kidneys shifted rostrally more than the left in camel embryos. Malik and Vais (1998) also reported that the right and left kidneys shifted rostral with advancement of age in ruminants. This cranial positioning of the kidneys might be due to relative variation in growth of different organs in the abdominal and pelvic cavities during various stages of embryonic development (Salehi and Morovati, 2012). Adipose tissue was found at the hilus of each kidney. Dyce, 1995 reported similar finding in ruminant, the fat sometime enough to

hide the kidney completely. The fat protects against distorting pressures from neighbouring organs. Salehi et al., (2012) also reported that adipose tissue surrounded the hilus and sides of the kidney.

A distinct renal pelvis with a well developed ureter originating from the middle of the renal pelvis was found in all kidneys, similar to Salehi et al., (2012) and Sarma et al., (2007). This confirms the development of renal pelvis and ureter in early phase of the gestation period in *Camelus dromedarius* embryos (Salehi and Morovati, 2012). The right ureter was found to have higher length than the left one in all foetuses. Abdalla et al., (1974) also reported that the right ureter of the camel is longer than the left considering the positions of the kidneys and the bladder in adult kidneys.

The values of foetal weight were found to increase significantly with advancement of gestational age. This is in accordance with Sonfada et al., (2012) who observed that there was an increase in body weight across the trimesters in the foetus with advancement in pregnancy, and Anderson et al., (1987) that there were obvious body weight changes in MDX mice which seem to increase with age.

The weights of left and right kidneys were found to be equal in the first trimester foetuses. Though there was no significant difference observed between the weight of the right and left kidneys in the study, there was significant difference between the left and right kidneys in both length and width in the study. The length and width values of the right kidneys were found to be greater than the left ones. The volume of the left kidneys was greater than the right one in first trimester. This finding is in contrast to the work of Salehi et al., (2012).

In the second and third trimesters such values of all parameters measured were found have greater values in the left kidneys than in the right ones. The lengths of the left kidneys were found to have higher values than their right counterpart in the 2nd and 3rd trimesters. Similar observations were made for the width values. The weight values of the left kidney were found significantly higher than that of the right one in the 2nd and 3rd trimester foetuses. This was in accordance with the findings of Salehi et al., (2012) in camel embryos and that of Dowelmadina, 2012 in adult camels and that of Sarma et al., 2007 in crossbred pig foetuses. Al-Ani (2004) reported that the larger kidney which was twice that in cattle and four times that of sheep was possibly due to the adaptation of camel to arid desert life. Constantinesu, (2004) reported that the greatest dimension of kidney may be due to size of the animal, direction and position.

Histological development of the renal tissues was also found to be variable across the three trimesters. The camel kidney present two distinguishing zones i.e cortex and medulla at first, second and third trimesters which is similar to the finding of Franco et al., (2004) on llama, guanaco and on goat. This is however in line with varying number of glomerulus and different levels of development of the duct system in the cortex and medulla of the camel kidneys at different trimesters. Stages of growth observe ranges from the level of mesenchyme cells, connective tissue proliferation, developing glomerulus, developed glomerulus, immature developing duct system and finally to matured duct system in the medulla and cortex of the kidneys. This is in line with that observed by franco et al., (2004) on Llama and the observations of El-Salmi and Amri (2012) on Sudanese sheep. Histological sections showed the presence of thick capsule which becomes thicker with advancement of gestational age was noticed in the kidneys.

The cortex of the first trimester kidney was found to have numerous mesenchymal cells and connective tissues with few developing glomerulus. This is in agreement with Onarlioglu et al., (1997) that observed no developed glomeruli in the kidneys where dense mesenchymal tissue take place in early stage of development in rat foetus. The number of mesenchyme cells reduced in the second trimester, together with few fully developed glomerulus and immature developing duct system.

The camel is the only domestic animal that has collecting duct system in the cortex. Numerous developed glomerulus and matured duct system with moderate developing glomerulus and connective tissue were seen at the third trimester. Extensive coiling of the tubular duct system was evident in the third trimester kidneys. This is in line with that observed by Franco et al., (2004) on Llama and El-Salmi and Amri (2012) on Sudanese sheep.

The medulla of the first trimester kidney was found to have various types of connective tissue (elastic, collagen and reticular) fibers and mesenchyme cells. Numerous developing collecting ducts, few developed collecting ducts and few mesenchyme cells were seen in the second trimester. This is also in line with that observed by Franco et al., (2004) on Llama and El-Salmi and Amri (2012) on Sudanese sheep. At third trimester the kidneys showed predominant developed collecting ducts and few mesenchyme cells, which showed the kidneys, were still developing. This trend of growth was in accordance with the findings of Patten and Carlson (1977), who reported that variable growth and structural diversities at different stages of development of an organ is a normal phenomenon for accommodating and molding of the organ.

5. Conclusion

In conclusion, the histological differentiation of the camel kidney is similar to that of sheep, goat, horse, llama, and guanaco but differs from that of the cat, dog and pig. Mean while, considering the histological features observed in the study, the camel kidney may become fully matured and functional at post-natal age.

Acknowledgement

I wish to show my sincere gratitude to Mr. M. Jimoh and Mr. O. Shola of the department of veterinary Anatomy, Faculty of veterinary medicine, Usmanu Danfodiyo university sokoto, for a job well-done in preparing the histological slide.

References

- Abdalla, M.A., 1973. Anatomical study of the urinary system of the camel (*Camelus dromedarius*). M. V. Sc. Thesis, University of Khartoum.
- Abdalla, M.A., Abdalla, O., 1979. Morphometric observations on the kidney of the camel, *Camelus dromedarius*. J. Anat, 129: 45-50.
- Abubakar, U.B., Mohammed, F.U., Shehu, S. A., Mustapha, R.A., 2010. Foetal Wastage in Camels Slaughtered (*Camelus dromedarius*) at Maiduguri Abattoir, Borno State Nigeria. Int. J. Trop. Med. 5(4), 86-88.
- Almathen, F.S., Mwacharo, J., Hanotte, O., 2012. Genetic Diversity and Relationships of Indigenous Saudi Arabia Camel (*Camelus dromedarius*) Populations. In: The proceedings of the Proceedings of the 3rd Conference of the International Society of Camelid Research and Development, Pp 40.
- Aly, K.H., 2007. Development of the mesonephros in camel (*Camelus dromedarius*). Anatomia, Histologia, Embryologia, 36, 58-61.
- Anderson, J.E., Ovalle, W.K., Bressler, B.H., 1987. Elctron microscope and audioradiograph characterization of hind limb muscle regeneration of MDX Mouse. Anat. Rec., 23(11), 243-257.
- Bello, A., Onyeausi, B.I., Sonfada, M.L., Adeyanju, J.B., Umar, A.A., Umaru, M.A., Shehu, S.A., Hena, S.A., 2012. Histomorphological studies of the prenatal development of oesophagus of one humped camel (*camelus dromedarius*): Sci. J. Agr., 1(4), 100-104.
- Bello, A., Onyeausi, B.I., Sonfada, M.L., Adeyanju, J.B., Umaru, M.A., 2012. A biometric study of the digestive tract of one-humped camel (*camelus dromedarius*) fetuses. Sci. J. Zool., 1(1), 11-16.
- Bello, M.B., Garba, H.S., Sonfada, M.L., 2008. Foetal Wastage in camels slaughtered at Sokoto Munincipal abattoir. Sokoto J. Vet. Sci., 7(1), 46-49.
- Buhari, B.K., 2008. Sokoto state investment promotion committee. Trade.
- Dowelmadina, I.M.M., 2012. Anatomical Characteristics of kidney in one-humped camel (*Camelus dromedarius*) in Sudan. In: The proceedings of the 3rd Conference of the International Society of Camelid Research and Development, pp 340-342.
- Dyce, K.M., Sack, W.O., Wesing, C.J.G., 1996. Textbook of Veterinary Anatomy 2nd edition, W.B. Saunders Company Philadelphia Pennsylvania, 19106 Pp175-179.
- Elwishy, A.B., Hemeida, N.A., Umar, M.A., Mobarak, A.M., El-Sayed, M.A.I., 1981. Changes in the pregnant camel with special reference to Foetal growth. British Vet. J., 37, 527-537.
- Eurell, J.A., Frappier, B.L., Dellmann, H., 2006. Textbook of Veterinary Histology. 6th Edition Lea and Febiger, Philadelphia, pp 212-219.
- Fayed, R.H., 2008. Adaptation of the Camel to Desert environment.
- Franco, A., Masot, A.J., Gómez, L., Redondo, E., 2004a. Morphometric and immuno histochemical study of the rumen of red deer during prenatal development. J. Anatom., 204, 501-513.
- Kataria, N., Kataria, A.K., Kumar, V.A., Garg, S.L., Singh, M.S., 2007. Solute loads and transfer function of kidney in dromedary camel during dehydration and rehydration in winter and summer. Vet. Arhiv., 77(3), 237-246.
- König, H.E., Liebich, H.G., 2009. Veterinary Anatomy of Domestic Mammals Textbook and colour Atlas 4th revised edition schattauer stuttgart New York pp 393-400.
- Kuure, S., Vuolteenaho, R., Vainio, S., 2000. Kidney morphogenesis: cellular and molecular regulation. Mech. Develop., 92, 31-45.

- McGeady, T.A., Quinn, P.J., FitzPatrick, E.S., Ryan, M.T., 2006. *Veterinary Embryology*. Blackwell Publishing, UK, ISBN: 9781405111478, pp 233-240
- Onarlioglu, B., Gursoy E., kaloglu, C., 1997. Renal Morphogenesis of rat fetus. *Turk. J. Anim. Sci.* 21, 493-496.
- Ouajd, S., Kamel, B., 2009. Physiological Particularities of Dromedary (*Camelus dromedarius*) and Experimental Implications. *Scand. J. Lab. Anim. Sci.*, 36(1).
- Salehi, E., Morovati, M.S., 2012. Kidney Morphogenesis during Prenatal Development in *Camelus dromedaries* Embryoes. *J. Anim. Vet. Adv.*, 1(6), 822-825.
- Sarma, K., Ahmed, B., 2007. Morphogenesis of kidney in crossbred pig (*Sus scrofa*) foeti. *Int. J. Morphol.*, 25(1), 129-132.
- Septimus, S., James, D.G., 1975. *The Anatomy of the domestic animals*. 5th Edition. vol.1 Pp 21-22. Pub. philadelphia: Saunder's company, Pp 564-579.
- Shehu, S.A., Ojo, S.A., Hambolu, J.O., Ibrahim, N.D.G., Hena, S.A., 2012. External features (Developmental Horizons) in developing camel foetus as a guide to Age estimation. *J. Vet. Adv.*, 2(4), 178-183.
- Skidmore, J.A., 2005. Reproduction in dromedary camels: An update. *Anim. Reprod.*, (2)3, 161-171.
- Smuts, M.M.S., Bezuidenhout, A.J., 1987. *Anatomy of the Dromedary* Oxford Clarendon Press, 133-134.
- Sonfada, M.L., 2008. Age related changes in musculoskeletal Tissues of one-humped camel (*Camelus dromedarius*) from foetal period to two years old. A Ph.D Thesis, Department of Veterinary Anatomy, Faculty of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto, Nigeria.
- Sonfada, M.L., Kwari, H.D., Umar, A.A., Shehu, A.A., Wiam, I.M., Hena, S.A., Danmaigoro, A., Onyeausi, B.I., 2012. Fetal Age Estimation in Dromedarian Camel Using Developmental Horizons. In: The proceedings of the Proceedings of the 3rd Conference of the International Society of Camelid Research and Development, Pp115-117.
- Sperber, I., 1944. Studies on the mammalian kidney. *Zoologiska Bidrag fran Uppsala* 22, 249-432.
- Tayeb, M.A.F., 1948. Urinary system of the camel. *J. Ameri. Vet. Med. Assoc.*, 113, 568 -572.
- Tisher, C.C., 1971. Relationship between renal structure and concentrating ability in the rhesus monkey. *Ameri. J. Phy.*, 220, 1100-1106.
- Tisher, C.C., Scchrier, R.W., McNeil, J.S., 1972. Nature of urine concentrating mechanism in the macaque monkey. *Ameri. J. Phy.*, 223, 1128-1137.
- Vimtrup, B.J., Schmdt-Nielsen, B., 1952. The histology of the kidney of the Kangaroo rats. *Anatom. Rec.*, 114, 515-528.
- Weibel, E.R., Gomez, D.M., 1962. A principle for counting tissue structures on random sections. *J. Appl. Phy.*, 17, 343-348.
- Weibel, E.R., 1963 a. Principles and methods for the morphometric study of the lung and other organs. *Lab. Invest.*, 12, 131-155.
- Weiber, E.R., 1963b. *Morphometry of the Human Lung*. Berlin: Springer Verlag.
- Wirz, H., 1954. The production of hypertonic urine by the mammalian kidney. In *The Kidney*, Ciba Foundation Symposium, pp. 38-49. Boston: Little, Brown and Co.
- Wirz, H., Dirix, R., 1973. Urinary concentration and dilution. In *Handbook of Physiology*, (section 8th ed. J. Orloff, R. W. Berliner & S. R. Geiger). Washington D.C.: American Physiological Society.